



www.BioOne.org

## 海外の事例

# BioOne

~ 全世界540機関、550万人が利用する

生物系電子ジャーナル・コレクション~

BioOne 指定代理店（北米地区を除く）

CSA Japan 沢田さやか



## BioOneとは・・・

www.BioOne.org

BioOneは、  
学会を含む非営利組織が刊行する  
質の高い科学誌に対するコスト効率のよい  
アクセスを確保・維持するために  
共同で尽力することをその使命とする。



## BioOneとは・・・

www.BioOne.org

- 学術図書館と非営利出版社による真の合併事業
- 初期財源の大部分が大学図書館からの寄付
- 図書館、出版社が技術的・財政的に持続可能な電子出版事業を支援



## BioOneとは・・・

www.BioOne.org

### BioOne後援団体

- 米国生物科学学会 (AIBS)
- SPARC (Scholarly Publishing and Academic Resources Coalition)
- ARL (Association of Research Libraries)
- The University of Kansas
- The Greater Western Library Alliance
- Allen Press, Inc.



## BioOneとは・・・

www.BioOne.org

- 理事会 図書館界：出版界 (50%:50%)
- 非営利目的でありながら費用効果(コスト回収など)の高いビジネスモデルの確立と実践に最大限の努力



## BioOneとは・・・

www.BioOne.org

2004年7月現在

生物学、植物学、生態学、環境科学系ジャーナルが中心

- ジャーナル - 70
  - 出版社 - 55
  - 電子書籍 - 1
  - 論文 - 27,863
  - OA ジャーナル - 3
- 巻数- 662  
ページ数- 217,032  
著者数 - 73,618  
号数- 1,687



## BioOneとは・・・

www.BioOne.org

- インパクトファクターの高いピア・レビュー誌を集めた電子ジャーナル・コレクション
- フルテキスト形式: PDF・SGML
- PubMed, Medline, NBII, JSTOR, Cross-Ref, BIOSIS, CSA等とのリンク対応



# BioOne検索画面

www.BioOne.org

HTML and PDF of full text of all articles – fully searchable

Select Journal(s)

All Journals
AMBIO: A Journal of the Human Environment
The American Biology Teacher
American Fern Journal
The American Midland Naturalist

Select Date Range

	Year	Month
From:	All	All
To:	All	All

Search (Note: Boolean operators AND, OR, NOT will be ignored inside these boxes.)

Title	with all the words	
Abstract	with all the words	
Full-Text	with all the words	
References	with all the words	
Author First Name	with all the words	
Author Last Name	with all the words	

Show  closest matches. Sort by






HOME

BiologyBrowser  
FROM BIOSIS

Soon you'll see it as irreplaceable.

> About > Search > Browse > Publishers > Subscribers > Help > FAQ



Full Text View

[\[PDF\]](#) [\[Create Reference\]](#)

## TABLE OF CONTENTS

[\[Material and...\]](#) [\[Results and...\]](#) [\[References\]](#) [\[Tables\]](#) [\[Figures\]](#)

---

*The Coleopterists Bulletin*: Vol. 58, No. 1, pp. 37–43.

### New Open Aquarium System to Breed Larvae of Water Beetles (Coleoptera: Dytiscidae)

Toshio Inoda,<sup>a</sup> and Shinji Kamimura<sup>a</sup>

<sup>a</sup>*Department of Life Sciences, Graduate School of Arts and Sciences, The University of Tokyo, Komaba 3-8-1, Meguro, Tokyo 153-8902, JAPAN. E-mail: [kam@nano.c.u-tokyo.ac.jp](mailto:kam@nano.c.u-tokyo.ac.jp)*

#### ABSTRACT

For conservation purposes and to supply rare insects for laboratory use, a system for artificial breeding is crucial. However, in the case of carnivorous freshwater insects such as diving beetles, constant conditions in aquariums are difficult to maintain due to their high rate of food consumption. Furthermore, surface rippling caused by the pumping system for water circulation hinders the respiration of small larvae. We developed a new open aquarium system without water circulation that was successfully applied to the rearing of larvae of diving beetles, *Dytiscus sharpi* (Wehncke) (Coleoptera: Dytiscidae). In comparison to conventional methods, a high proportion of larvae developed into adult insects. The size of reared adults was almost the same as those of field-collected adults. The new method could be applied to the conservation and breeding of other rare species, such as water beetles and water bugs.

Manuscript Received by the Society 2 May 2002

Manuscript Accepted 30 April 2003

Publication date 2 April 2004

The carnivorous larvae of *Dytiscus* species prey on other animals, e.g., insect larvae, amphibian larvae, isopods, or fish fry (Blunck 1923; Balduf 1935; Wernberg-Lund 1943; Jeffries 1988; Johanson and Nilsson 1992). They first inject digestive enzymes through the mandibles into the body of the prey then

## Adult Insects

Wild diving beetles (*Dytiscus sharpi*, Fig. 1D) were collected (n = 27) at Boso Peninsula in Chiba, Japan, and were mated under the

They were kept outdoors in plastic tanks (74 × 39 × 40 cm<sup>3</sup>) with 10 cm water depth. They were fed small pieces of dried sardines once a week. De-chlorinated water was automatically supplied every 6h by overflowing the tanks. Japanese parsley (*Oenanthe javanica* (Blume)) was placed in the tanks to provide resting and oviposition substrates. Usually, mating behavior started in November and eggs were found in March after wintering (Inoda 2003).

## Juveniles

Thirty newly hatched first instars (Fig. 1A) were collected and kept isolated in each tank of a new open aquarium system (details below). To obtain second instars for the experiments, the remaining first instars were isolated in plastic containers (diameter 4 cm × height 7 cm, 5 cm water depth). They were fed *Aseilus hilgendorfi* (Bovalius) (isopods) and/or tadpoles of *Rana ornativentris* (Werner). Fresh de-chlorinated tap water was supplied after feeding. Following development to second instars (Fig. 1B), they were transferred into the tanks of the open aquarium system (n = 94). Some were kept in the same tank partitioned by a 3mm mesh plastic net (Fig. 3). One hundred-150 tadpoles of *R. ornativentris* were supplied to each tank every 7–10 days.

Third instar larvae (Fig. 1C) required no food prior to pupation and were transferred into plastic containers (11 × 8 × 7 cm<sup>3</sup>) filled with moist soil and kept indoors at 10° to 24°C until emergence. All other stages were kept outdoors (8°–21°C).

## Aquarium System

The open aquarium system used in the present study consisted of three compartments; a water reservoir, a filtering unit and a tank for breeding. A schematic drawing and photographs of the system are shown in Figure 2.

Between the water reservoir (100 liter) and the filtering unit, a submersible pump (EHEIM 1046, 300 liter/h) was placed and run for 20 min every 6h. While the pump was working, the water level in the reservoir tank was kept constant by supplying fresh tap water through a shut-off valve connected to a float. The filtering unit consisted of connected columns of a pre-filter for crude filtering (5µm, Organo, Tokyo) and a carbon filter for de-chlorination (Organo, Tokyo). Our preliminary observation indicated that any residual chlorine caused irreversible damage both to larvae of *D. sharpi* and tadpoles of *R. ornativentris*.

Through a silicon tube (inner diameter 4 mm), de-chlorinated water was then gently introduced into the breeding tanks (30 × 20 × 10 cm<sup>3</sup>, 7 cm water depth) that were placed in stacks as shown in Figure 2. Water overflowing from the upper tanks supplied the lower ones. Each tank was equipped with outlet tubes of vinyl chloride (inner diameter 13 mm) and rubber gaskets (inner diameter 17.8 mm). When we kept two or three larvae in the same tank, plastic nets were used for partitioning in order to prevent cannibalism (Fig. 3).

## Measurement of Body Size

To investigate whether the present experimental conditions were appropriate for breeding, we compared the sizes of reared adults with those of wild ones. Dorsal aspects of adult insects were photographed with a digital camera (CoolPix 990, Nikon, Tokyo) and the length of body, elytra, pronotum, and maximal body width were determined. Adult insects were also weighted. A non-paired *t*-test was used to test the statistical significance of differences between mean



[sharpi](#), which has so far been obscure.

### Acknowledgments

We thank H. Katakura (Department of Life Sciences, University of Hokkaido) and M. Ohara (Systematic Entomology, Hokkaido University Museum) for the critical reading of the manuscript.

### Literature Cited [Return to TOC](#)

Balduf, W. V. 1935, The bionomics of entomophagous Coleoptera: Reprint 1969 by E. W. Classey Ltd., Hampton, England. 200 pp.

Blunck, H. 1923, Die Entwicklung des [Dytiscus marginalis](#) L. vom Ei bis zur Imago. 2. Teil. Die Metamorphose (B. Das Larven- und das Puppenleben): Zeitschrift für Wissenschaftliche Zoologie. 121 117–391. [Links ▶](#)

Brosher, F. 1913, Recherches sur la respiration des insectes aquatiques: Etude anatomique et physiologique du système respiratoire chez les larves du genre [Dytiscus](#). Annales de Biologie Lacustre, T. 6, 120–147. Bruxelles.

Inoda, T. 2003, Mating and reproduction of predaceous diving beetles, [Dytiscus sharpi](#), observed under artificial breeding conditions: Zoological Science. 20 377–382. [Links ▶](#)

Jeffries, M. 1988, Individual vulnerability to predation: the effect of alternative prey types: Freshwater Biology. 19 49–56. [Links ▶](#)

Johansson, A., and A. N. Nilsson. 1992, [Dytiscus latissimus](#) and [D. circumcinctus](#) (Coleoptera, Dytiscidae) larvae as predators on three case-making caddis larvae: Hydrobiologia. 248 201–213. [Links ▶](#)

Linnaeus, C. 1758, Systema naturae per regna tria naturae secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis: Tomus. Editio decima, reformata. Stockholm. 823 + 1 pp. [Photographic facsimile, British Museum (Natural History), London&rsqb.

Ministry of the Environment, Government of Japan. 2000, Red data list of Japan. (in Japanese):

Régimbart, M. 1899, Revision des Dytiscidae de la région Indo-Sino-Malaise: Annales de la Société Entomologique de France. 68 186–367. [Links ▶](#)

Sharp, D. 1884, The water-beetles of Japan: Transactions of the Entomological Society of London, 1884, 439–464.

Wehncke, E. 1875, [Dytiscus sharpi](#) n. sp: Stettiner Entomologische Zeitung, 1875, 500.

Wesenberg-Lund, C. 1943, Biologie der Susswasserinsekten: Nordisk forlag, Kopenhagen. 682 pp.

Wilson, C. B. 1923, Water beetles in relation to pondfish culture, with life histories of those found in fishponds at fairport, Iowa: Bulletin U.S. Bureau of Fisheries. 39 231–345.

[Go to Print Version](#)

Results of Search in every Kingdom for all exactly for 'Oenanthe javanica'

### ***Oenanthe javanica* (Blume) DC.**

Taxonomic Serial No.: 506796

#### Taxonomy and Nomenclature

Kingdom: Plantae  
 Taxonomic Rank: Species  
 Synonym(s):  
 Common Name(s): Java waterdropwort

**Taxonomic Status:**

Current Standing: accepted

**Data Quality Indicators:**

Record Credibility Rating: verified - standards met

#### Taxonomic Hierarchy

Kingdom	<a href="#">Plantae</a> -- Planta, plantes, plants, Vegetal
Subkingdom	<a href="#">Tracheobionta</a> -- vascular plants
Division	<a href="#">Magnoliophyta</a> -- angiospermes, angiosperms, flowering plants, phanérogames, plantes à fleurs, plantes à fruits
Class	<a href="#">Magnoliopsida</a> -- dicots, dicotylédones, dicotyledons
Subclass	<a href="#">Rosidae</a>
Order	<a href="#">Apiales</a>
Family	<a href="#">Apiaceae</a>
Genus	<a href="#">Oenanthe</a> L. -- waterdropwort
Species	<i>Oenanthe javanica</i> (Blume) DC. -- Java waterdropwort

**Direct Children:**

Subspecies: [Oenanthe javanica ssp. javanica](#) (Blume) DC. -- Java waterdropwort  
 Subspecies: [Oenanthe javanica ssp. stolonifera](#) (Wallich ex DC.) Murata -- stolon waterdropwort

#### References

**Expert(s):**

Expert: Mark W. Skinner


**Tables** [Return to TOC](#)

**Table 1.** Comparison of body size between reared and wild adult insects

Sex		Body length (mm)	Elytral length (mm)	Pronotum length (mm)	Body width (mm)	Weight (g)
Male	Wild (n = 13)	31 ± 1.7	24 ± 1.2	5.2 ± 0.4	16 ± 0.8	1.8 ± 0.2
	Bred (n = 38)	31 ± 1.2	23 ± 1.2	5.1 ± 0.4	16 ± 0.7	1.9 ± 0.3
Female	Wild (n = 14)	30 ± 1.9	22 ± 0.9	4.6 ± 0.6	15 ± 0.5	1.6 ± 0.1
	Bred (n = 29)	30 ± 1.5	22 ± 1.0	4.8 ± 0.5	15 ± 0.9	1.7 ± 0.2

Mean values ± SD.

**Figures** [Return to TOC](#)



Click on thumbnail for full-sized image.


**Fig. 1.** Photographs showing each stage of larvae of *D. sharpi* kept in the open aquarium system. **A)** First instar larva 5 days after hatching; **B)** second instar larva 10 days after hatching; **C)** third instar larva 32 days after hatching; **D)** a reared adult. Bars indicate 10 mm.



# 利用率の拡大

www.BioOne.org

2003年冬に1ヶ月の利用者数100万人を突破！





## 拡大する収録誌

www.BioOne.org

- ジャーナル数
  - 4年間でジャーナル数が2倍に！
    - 2001年 - 35 誌
    - 2004年 - 70 誌
  - 2004年末までに3誌、来年以降もさらに追加予定
- コレクションの排他性
  - 90%のジャーナルはBioOneのみで提供  
(他プラットフォームで提供されているのは4誌のみ)



## 高い顧客満足度

www.BioOne.org

- 高い更新率 - 99.9%
- 経済状況の変化にも関わらず、新規導入数は順調に増加
- 北米の図書館は、質の高い学術情報が合理的な価格で流通し、またそれらが世界中に広まるようBioOneを継続支援





## BioOneとは・・・まとめ

www.BioOne.org

- **高品質/低価格商品**
  - ピア・レビュー誌
  - ISI rankedジャーナル
- **お値ごろ商品**
  - 過去4年間の価格上昇率16%
  - フォーカス・グループ調査の結果
    - 値上がりが十分でない！
    - コスト効率の良い商品により投資していきたい。



# お問い合わせ

www.BioOne.org

BioOneに関するご質問、お問い合わせは、

CSAジャパン

[ssawada@csa.com](mailto:ssawada@csa.com)

045-222-8573

までご連絡ください。